

# 1016-351-01

## Probability

### Problem Set 4

Assigned 2012 January 10  
Due 2012 January 17

Show your work on all problems! If you use a computer to assist with numerical computations, turn in your source code as well.

- 1 Devore Chapter 3, Problem 74
- 2 Devore Chapter 3, Problem 76
- 3 Devore Chapter 3, Problem 86
- 4 Devore Chapter 3, Problem 88
- 5 Computational Exercise (Extra Credit)

The hypergeometric distribution

$$h(x; n, M, N) = \frac{\binom{M}{x} \binom{N-M}{n-x}}{\binom{N}{n}} \quad (5.1)$$

can be approximated by a binomial distribution

$$b(x; n, p) = \binom{n}{x} p^x (1-p)^{n-x} \quad (5.2)$$

with  $p = M/N$ , when  $M$ ,  $N$ , and  $N - M$  are all large.

- a. Using a computer, plot the pmfs  $h(x; 10, 12, 20)$  and  $b(x; 10, .6)$  over the range of possible  $x$  values. (Recall that if you use matplotlib, the binomial coefficient can be imported from scipy with `from scipy import comb`.)
- b. Using a computer, plot the pmfs  $h(x; 10, 120, 200)$  and  $b(x; 10, .6)$  over the range of possible  $x$  values.
- c. Another large-number approximation is that the binomial distribution tends towards the Poisson distribution

$$p(x; \mu) = \frac{e^{-\mu} \mu^x}{x!} . \quad (5.3)$$

On the same set of axes, plot the Poisson pmf  $p(x; 3)$  and the binomial pmfs  $b(x; 12, .25)$  and  $b(x; 300, .01)$ , for  $x$  between 0 and 12, inclusive. (If you use matplotlib, you can also import a factorial function from scipy with `from scipy import factorial`.)