

MATH 252-01: Probability and Statistics II

Problem Set 2

Assigned 2019 January 22

Due 2019 January 29

Show your work on all problems! If you use a computer to assist with numerical computations, turn in your source code as well.

1 Devore Chapter 7, Problem 4

2 Devore Chapter 7, Problem 22

3 Devore Chapter 7, Problem 34

4 Computational Exercise: Bootstrapping

The *Bootstrap* technique provides a model-free way to estimate the error associated with a point estimate from a reasonable-sized sample. If the sample values are $\{x_i\} = x_1, x_2, \dots, x_n$, we create a total of B bootstrap samples $\{x_i^{(1)}\}, \{x_i^{(2)}\}, \dots, \{x_i^{(B)}\}$. Each sample has size n , and each is generated by drawing *with* replacement from $\{x_i\}$. (So in general, a given bootstrap sample will have some repeated values, and some values not represented.) To estimate the error associated with the sample mean $\bar{x} = \frac{1}{n} \sum_{i=1}^n x_i$ of the original sample, we calculate the means of the B bootstrap samples, $\bar{x}^{(1)} = \frac{1}{n} \sum_{i=1}^n x_i^{(1)}$, $\bar{x}^{(2)} = \frac{1}{n} \sum_{i=1}^n x_i^{(2)}$, \dots , $\bar{x}^{(B)} = \frac{1}{n} \sum_{i=1}^n x_i^{(B)}$, and then take the (sample) variance of $\bar{x}^{(1)}, \bar{x}^{(2)}, \dots, \bar{x}^{(B)}$,

$$s_{\bar{x}}^2 = \frac{1}{B-1} \sum_{j=1}^B (\bar{x}^{(j)} - \bar{\bar{x}}^*)^2$$

where $\bar{\bar{x}}^* = \frac{1}{B} \sum_{j=1}^B \bar{x}^{(j)}$ is the average of the means of the bootstrap samples. The bootstrap estimate of the error associated with the original sample mean \bar{x} is then $\sqrt{s_{\bar{x}}^2}$.

- a. Download the following data set which is a sample of size $n = 5$
http://ccrg.rit.edu/~whelan/courses/2019_1sp_MATH_252/data/ps02_prob4_small1.dat
using the username and password given in class; generate $B = 8$ bootstrap samples by randomly choosing 8 sets of 5 values each from the original dataset. Be sure to turn in a table containing these 8 bootstrap samples.
- b. Calculate the mean of each of your 8 bootstrap samples.

- c. Calculate the bootstrap error as the sample standard deviation of this set of 8 bootstrap means. Note that this is not a robust use of the bootstrap method, since we have a small number of samples, but it's a way to see how the calculation works explicitly.
- d. Repeat the calculation using the data set http://ccrg.rit.edu/~whelan/courses/2019_1sp_MATH_252/data/ps01_prob4.dat which has size $n = 121$, with $B = 200$ bootstrap samples. (You don't need to print out the full set of 200×121 values, just calculate the bootstrap error estimate and document the procedures.) If you have trouble automatically generating the bootstrap samples, you may use the datafile http://ccrg.rit.edu/~whelan/courses/2019_1sp_MATH_252/data/ps02_prob4_resampled.dat which contains 200 columns, each with a 121-row bootstrap sample.
- e. The data were actually generated from a $\text{Gamma}(1.5, 15)$ distribution; use this to calculate the standard error $\sqrt{V(\bar{X})}$. (*Hint*: the sample mean of a sample of size n from a $\text{Gamma}(\alpha, \beta)$ distribution is a statistic which follows a $\text{Gamma}(n\alpha, \beta/n)$ distribution.)